CoxHealth at Home Dermatology (drugs S)

					FAX: 1-417-269	-0092
Patient Informa				Shipping Informat		
Patient name: DOB:						
			NPI:			
		_ ❑kg ❑lbs Ht:❑cm ❑in				
Address:	ddress:				State: Zip:	
Apt/Suite: City: State: Zip:			Contact:			
Phone: Alternate:			Phone:	Phone: Alternate:		
Caregiver name: Relation:			Fax:			
	Local pharmacy: Phone:					
	nsurance plan: Plan ID:		If shipping to prescriber: D First Fill D Always D Never			
Please fax a copy of front and back of the insurance card(s).						
Clinical Information (Please fax all pertinent clinical and lab information)						
Diagnosis : L40.0 (Psoriasis vulgaris/Plaque Psoriasis/Nummular Psoriasis) L40.8 (Other psoriasis)						
□ L40.9 (Psoriasis, unspecified) □ L40.5 (Psoriatic arthritis) □ L73.2 (Hidradenitis Suppurativa) □ Diagnosis Date: TB test: □ Yes □ No Neg. Test Date: HBV: □ Yes □ No If yes, currently treated: □ Yes □ No						
RSA affected (%):	Affect	ed areas: \Box Palms \Box Soles \Box F	lead 🖸 Neck 🔲	Genitalia	Sufferiliy treateu. 🖬 res	
		Reason for Discontinuation of Th		Approximate Start D	Date Approximate End	Date
Prior Therapy			петару			Date
Comorbidities:					L	
Concomitant Medications:						
Allergies: NKDA Other:						
Has the patient received their starter dose(s)/kit? Yes; Start Date No						
Prescription				Quantity		Refill
		See forms A-E, F-R, and T-Z	Z for their respecitve	.		
	🗖 Inject 210 mg SC	on weeks 0, 1, and 2 followed by 2	210 mg			
^I Siliq [™] (brodalumab)	every 2 weeks thereafter			4 x 210 mg/1.5 mL	PFS	0
	□Inject 210 mg SQ every 2 weeks			2 x 210 mg/1.5 mL	PFS	- <u> </u>
Simponi [®] (golimumab) Psoriatic Arthritis	Inject 50 mg SQ once a month			1 x 50 mg/0.5 mL	SmartJect Autoinjector	
					PFS	
Skyrizi (risankizumab- rzaa)	Starter: Inject 150mg SQ at week 0, 4, then every 12 weeks therafter			2 x 75mg/0.83 mL	PFS	
	□ Maintenance: Inject 150mg SQ every 12 weeks					
☐ Simponi Aria [®] (golimumab)	□Infuse mg (2 mg/kg x kg) IV over 30 min at weeks 0			x 50 mg/4 ml	Vials	0
				· · · · · · · · · · · · · · · ·		
	□ Infuse mg (2 mg/kg x kg) IV over 30 min at week 4			x 50 mg/4 ml	Vials	
	and every 8 weeks thereafter			X 00 mg/4 m	VIGIO	
Stelara [®] (ustekinumab) Adult	□ Inject 45 mg SQ on Day 1 (≤100 kg)			1 x 45 mg/0.5 mL		
	□Inject 90 mg SQ on Day 1 (>100 kg)			1 x 90 mg/1 mL	PFS	0
	□ Inject 45 mg SQ on Day 29 and every 12 weeks thereafter (≤100 kg)			1 x 45 mg/0.5 mL		<u> </u>
	□ Inject 90 mg SQ on Day 29 and every 12 weeks thereafter (>100 kg)			1 x 90 mg/1 mL	PFS	
				, ř		
Injection Training Provided by: Physician's Office CoxHealth at Home Other:						
Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:						
		Stamp signature not allow	ed, physician signat	•		
Prescriber's Signature:				Da	te:	-

I authorize CoxHealth at Home. and its representatives to act as an agent to initiate and execute the insurance prior authorization process for this prescription and any future fills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to CoxHealth at Home.

Confidentiality Statement: This message is intended only for the individual or entity to which it is addressed. It may contain information which may be proprietary and confidential. It may also contain privileged, confidential information which is addressed. It may contain information which may be proprietary and confidential. It may also contain privileged, confidential information which is information which it is addressed. It may contain information which may be proprietary and confidential. It may also contain privileged, confidential information which is information which is addressed. It may contain information which may be proprietary and confidential. It may also contain privileged, confidential information which is information of distributing the Health Insurance Portability and Accountability Act (HIPAA). If you are not the intended recipient, please note that you are strictly prohibited from disseminating or distributing this information. If you received this communication in error, please notify the sender immediately by calling 1 855-419-4663 to obtain instructions as to the proper destruction of the transmitted material. Thank you.